


Probability Trees and the Multiplication Rule

CS231
Dianna Xu

1

The multiplication rule

- Also called the product rule
- If there are n_1 ways to do task 1, and n_2 ways to do task 2
 - Then there are $n_1 \times n_2$ ways to do both tasks in sequence
 - We must make one choice AND a second choice



2

Product rule example

- Sample question
 - There are 18 MATH majors and 17 CS majors
 - How many ways are there to pick one math major **and** one CS major?
- Total is $17 \times 18 = 306$

3

Product rule example

- How many strings of 4 decimal digits...
- Do not contain the same digit twice?
 - We want to chose a digit, then another that is not the same, then another...
 - First digit: 10 possibilities
 - Second digit: 9 possibilities (all but first digit)
 - Third digit: 8 possibilities
 - Fourth digit: 7 possibilities
 - Total = $10 \times 9 \times 8 \times 7 = 5040$
- End with an even digit?
 - First three digits have 10 possibilities
 - Last digit has 5 possibilities
 - Total = $10 \times 10 \times 10 \times 5 = 5000$

4

When the product rule is difficult to apply

- President, treasurer and secretary are to be chosen among A, B, C, D. A can not be president and either C or D must be secretary.
- Naïve application of the product rule:
 - President: 3
 - Treasurer: 3
 - Secretary: 2
 - Total = 18

5

Tree diagrams

- We can use tree diagrams to enumerate the possible choices
- Once the tree is laid out, the result is the number of (valid) leaves

6

r-permutation Formula

- There are n ways to choose the first element
 - $n-1$ ways to choose the second
 - $n-2$ ways to choose the third
 - ...
 - $n-r+1$ ways to choose the r^{th} element
- By the product rule, that gives us:

$$P(n,r) = n(n-1)(n-2)\dots(n-r+1)$$

13

r-permutations example

- How many ways are there for 3 students in this class to sit together?
- There are 50 students in the class
 - $P(50,3) = 50 \times 49 \times 48 = 117,600$
 - Note that the positions they take do matter

14

Permutations vs. r-permutations

- r-permutations: Choosing an ordered 5 card hand is $P(52,5)$
 - When people say “permutations”, they almost always mean r-permutations
 - But the name can refer to both
- Permutations: Choosing an order for all 52 cards is $P(52,52) = 52!$
 - Thus, $P(n,n) = n!$

15

Sample question

- How many permutations of $\{a, b, c, d, e, f, g\}$ end with a?
 - Note that the set has 7 elements
 - The last character must be a
 - The rest can be in any order
- Thus, we want a 6-permutation on the set $\{b, c, d, e, f, g\}$
- $P(6,6) = 6! = 720$
- Why is it not $P(7,6)$?

16