



CS 260 – Foundations of Data Science

Lecture 11 – Logistic Regression

10/19/2026

Adam Poliak



Announcements

HW5 due Tuesday night

Midterm 2 – November 19th

Midterm 1 – go over in lab today



Outline

Introduction to Logistic Regression

Cost function



Challenger Explosion Data



1	Date	Temperature	Damage Incident
2	04/12/1981	66	0
3	11/12/1981	70	1
4	3/22/82	69	0
5	6/27/82	80	NA
6	01/11/1982	68	0
7	04/04/1983	67	0
8	6/18/83	72	0
9	8/30/83	73	0
10	11/28/83	70	0
11	02/03/1984	57	1
⋮			
23	10/30/85	75	1
24	11/26/85	76	0
25	01/12/1986	58	1
26	1/28/86	31	Challenger Accident



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure

- 1) If you were forced to use linear regression for this problem, how could you encode y to make it real-valued?



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure

1) If you were forced to use linear regression for this problem, how could you encode y to make it real-valued?

You could choose stroke=0, drug overdose=1, epileptic seizure=2 (or some permutation)



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure

1) If you were forced to use linear regression for this problem, how could you encode y to make it real-valued?

You could choose stroke=0, drug overdose=1, epileptic seizure=2 (or some permutation)

2) What issues arise with making y real-valued?



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure

1) If you were forced to use linear regression for this problem, how could you encode y to make it real-valued?

You could choose stroke=0, drug overdose=1, epileptic seizure=2 (or some permutation)

2) What issues arise with making y real-valued?

Assumes some *ordering* of the outcomes that is probably not there!



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure

- 1) If you were forced to use linear regression for this problem, how could you encode y to make it real-valued?

You could choose stroke=0, drug overdose=1, epileptic seizure=2 (or some permutation)

- 2) What issues arise with making y real-valued?

Assumes some *ordering* of the outcomes that is probably not there!

- 3) What if you just had two outcomes (e.g. stroke and drug overdose) -- why is linear regression still not a good choice?



Why is linear regression a bad choice for classification?

Case Study: you need to identify the medical condition of a patient in the emergency room on the basis of their symptoms.

Possible conditions (y) are:

- Stroke
- Drug overdose
- Epileptic seizure

- 1) If you were forced to use linear regression for this problem, how could you encode y to make it real-valued?

You could choose stroke=0, drug overdose=1, epileptic seizure=2 (or some permutation)

- 2) What issues arise with making y real-valued?

Assumes some *ordering* of the outcomes that is probably not there!

- 3) What if you just had two outcomes (e.g. stroke and drug overdose) -- why is linear regression still not a good choice?



Desirata

Very positive

Zero

Very positive



Desirata

Very positive

Close to 0

Zero

0.5

Very positive

Close to 1



Desirata

Very positive

Close to 0

Zero

0.5

Very positive


Close to 1

Smooth transitions



Logistic (sigmoid) function

Transforms a continuous real number into a range of (0, 1)



Logistic Regression

- Binary classification $y \in \{0,1\}$
- Model will be

$$h_{\vec{w}}(\vec{x}) = \frac{1}{1+e^{-\vec{w} \cdot \vec{x}}}$$
$$= p(y = 1|\vec{x})$$

- Classification (already have $\vec{w}$)

$$\text{if } \vec{w} \cdot \vec{x} \geq 0 \Rightarrow \hat{y} = 1$$

$$\vec{w} \cdot \vec{x} < 0 \Rightarrow \hat{y} = 0$$



Logistic regression example

- If $p=1$ (one feature), can solve for x directly

$$w_0 + w_1 x \geq 0$$

$$w_1 x \geq -w_0$$

$$x \geq -\frac{w_0}{w_1}$$

- Ex: $\vec{w} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}$

$$x \leq \frac{3}{2} \text{ means predict } \hat{y} = 1$$



Outline

Introduction to Logistic Regression

Cost function & SGD



BRYN MAWR
COLLEGE



How to find $\vec{w}$?

- Need a cost function
- Can measure model performance with likelihood

$$L(\vec{w}) = \prod_{i=1}^n \underbrace{h_{\vec{w}}(\vec{x}_i)}_{\text{prob of } y=1}^{y_i} \underbrace{(1 - h_{\vec{w}}(\vec{x}_i))}_{\text{prob of } y=0}^{(1-y_i)}$$

want high



Cost function for logistic regression

$$J(\vec{w}) = \underbrace{-\log(L(\vec{w}))}_{\text{negative log-likelihood}}$$

↑
minimize

$$J(\vec{w}) = - \sum_{i=1}^n [y_i \log(h_{\vec{w}}(\vec{x}_i)) + (1 - y_i) \log(1 - h_{\vec{w}}(\vec{x}_i))]$$


- Single example $\vec{x}, y$

$$J(\vec{w}) = \begin{cases} -\log(h_{\vec{w}}(\vec{x})) & \text{if } y = 1 \\ -\log(1 - h_{\vec{w}}(\vec{x})) & \text{if } y = 0 \end{cases}$$



Single data point

$$J(\vec{w}) = \begin{cases} -\log(h_{\vec{w}}(\vec{x})) & \text{if } y = 1 \\ -\log(1 - h_{\vec{w}}(\vec{x})) & \text{if } y = 0 \end{cases}$$



Stochastic Gradient Descent for Logistic Regression (binary classification)

```
set  $\vec{w} = \vec{0}$ 
while cost  $J(\vec{w})$  is still changing:
    shuffle data points
    for  $i = 1, \dots, n$ :
         $\vec{w} \leftarrow \vec{w} - \alpha \nabla J_{\vec{x}_i}(\vec{w})$ 
    store  $J(\vec{w})$ 
```

$\underbrace{\hspace{10em}}$
derivative of $J(\vec{w})$ wrt
 x_i



3 important pieces to SGD

- Hypothesis function (prediction)

$$h_w(\mathbf{x}) = p(y = 1|\mathbf{x}) = \frac{1}{1 + e^{-w \cdot \mathbf{x}}}$$



3 important pieces to SGD

- Hypothesis function (prediction)

$$h_{\mathbf{w}}(\mathbf{x}) = p(y = 1|\mathbf{x}) = \frac{1}{1 + e^{-\mathbf{w} \cdot \mathbf{x}}}$$

- Cost function (want to minimize)

$$J(\mathbf{w}) = - \sum_{i=1}^n y_i \log h_{\mathbf{w}}(\mathbf{x}_i) + (1 - y_i) \log(1 - h_{\mathbf{w}}(\mathbf{x}_i))$$



3 important pieces to SGD

- Hypothesis function (prediction)

$$h_{\mathbf{w}}(\mathbf{x}) = p(y = 1|\mathbf{x}) = \frac{1}{1 + e^{-\mathbf{w} \cdot \mathbf{x}}}$$

- Cost function (want to minimize)

$$J(\mathbf{w}) = - \sum_{i=1}^n y_i \log h_{\mathbf{w}}(\mathbf{x}_i) + (1 - y_i) \log(1 - h_{\mathbf{w}}(\mathbf{x}_i))$$

- Gradient of cost wrt single data point $\mathbf{x}_i$

$$\nabla J_{\mathbf{x}_i}(\mathbf{w}) = (h_{\mathbf{w}}(\mathbf{x}_i) - y_i)\mathbf{x}_i$$



Stochastic Gradient Descent for Logistic Regression (binary classification)

```
set  $\vec{w} = \vec{0}$   
while cost  $J(\vec{w})$  is still changing:  
    shuffle data points  
    for  $i = 1, \dots, n$ :  
         $\vec{w} \leftarrow \vec{w} - \alpha \nabla J_{\vec{x}_i}(\vec{w})$   
    store  $J(\vec{w})$ 
```

$\underbrace{\hspace{10em}}$
derivative of $J(\vec{w})$ wrt
 x_i



Cost function

$$J(\vec{w}) = -[\underbrace{y}_{\text{probability distribution}} \log(\underbrace{h_{\vec{w}}(x)}_{\text{probability distribution}}) + \underbrace{(1-y)}_{\text{probability distribution}} \log(1 - h_{\vec{w}}(x))]$$

$$\underbrace{H(Y)}_{\text{entropy}} = - \sum_{y \in \text{vals}(Y)} p(y) \log p(y)$$

$$\underbrace{\text{Cross entropy}}_{\text{Cross entropy}} \quad H(p, q) = - \sum_{y \in \text{vals}(Y)} p(y) \log q(y)$$



Cost function as Cross Entropy

- Example
 - true: $y=0$, $1-y=1$
 - pred: $h=0.4$, $1-h=0.6$

$$H(\text{true}, \text{pred}) = -(0 \log(0.4) + 1 \log(0.6)) = 0.5$$



Handout



BRYN MAWR
COLLEGE



Handout

1. The output of logistic regression is a model that creates:
 - (a) a linear decision boundary
 - (b) a logistic decision boundary
 - (c) no decision boundary



Handout

1. The output of logistic regression is a model that creates:

- (a) a linear decision boundary
- (b) a logistic decision boundary
- (c) no decision boundary

2. We use logistic regression for:

- (a) classification
- (b) regression
- (c) both



Handout

1. The output of logistic regression is a model that creates:

- (a) a linear decision boundary
- (b) a logistic decision boundary
- (c) no decision boundary

2. We use logistic regression for:

- (a) classification
- (b) regression
- (c) both

3. Our hypothesis in logistic regression is:

$$h_{\mathbf{w}}(\mathbf{x}) = p(y = 1|\mathbf{x}) = \frac{1}{1 + e^{-\mathbf{w} \cdot \mathbf{x}}}$$

If $\mathbf{w}$ is the zero vector (as it would be when starting SGD), what is the probability $y = 1$?

Handout

1. The output of logistic regression is a model that creates:

- (a) a linear decision boundary
- (b) a logistic decision boundary
- (c) no decision boundary

2. We use logistic regression for:

- (a) classification
- (b) regression
- (c) both

3. Our hypothesis in logistic regression is:

$$h_{\mathbf{w}}(\mathbf{x}) = p(y = 1|\mathbf{x}) = \frac{1}{1 + e^{-\mathbf{w} \cdot \mathbf{x}}}$$

If $\mathbf{w}$ is the zero vector (as it would be when starting SGD), what is the probability $y = 1$?


$$1/2$$

Handout

Say I train a binary logistic regression model (i.e. outcomes $\in \{0, 1\}$) and end up with $\hat{\mathbf{w}} = [\hat{w}_0, \hat{w}_1]^T = [-4, -5]^T$. What is the decision boundary? Sketch a graph of this logistic model and label the decision boundary. How would you classify a new point $x_{\text{test}} = -2$?


$$\frac{1}{1 + e^{-(-4 - 5x)}}$$

predict $y=1$



Handout

5. The graph below shows the cost for logistic regression as a function of the hypothesis $h_w(\mathbf{x})$, for one example $\mathbf{x}$. Which curve corresponds to the true label $y = 0$ and which corresponds to $y = 1$?



Single example $\vec{x}, y$

$$J(\vec{w}) = \begin{cases} -\log(h_{\vec{w}}(\vec{x})) & \text{if } y = 1 \\ -\log(1 - h_{\vec{w}}(\vec{x})) & \text{if } y = 0 \end{cases}$$

